

PROVIDING ONLINE SELF-TESTS TO IDENTIFY OR ASSESS BEHAVIOR OR HEALTH DISORDERS RELATED TO CHRONIC STRESS AT WORK BY HEALTH PROFESSIONALS IN ANESTHESIOLOGY AND INTENSIVE CARE IN FRANCE

EAPH conference 2015 20th and 21st April 2015 - Barcelona, Spain Doctors' Resilience. Building European Networks through Research and Practice



DOPPIA M-A ¹, LIEUTAUD T ², ARZALIER-DARET S. ¹, MERTES P-M ³ on behalf of Members of SMART Committee of the French College of Anaesthetists and Intensivists / Committee for Anesthesiologists' Occupational Health)

> Address: CFAR - Collège Français des Anesthésistes Réanimateurs 74, rue Raynouard 75016 PARIS (France) - email : contact@cfar.org

Pôle Anesthésie Réanimation SAMU, CHU CAEN - Côte de Nacre - 14033 CAEN Cedex 09 - FRANCE

² Equipe TIGER, Centre de Recherche en Neurosciences, Lyon (France) UMRESTTE, UMR-T9405, IFSTTAR, Université Claude Bernard de Lyon ³ President of the CFAR, CHRU de Strasbourg Nouvel Hôpital Civil, Service d'Anesthésie-Réa Chirurgicale, 1 place de l'Hôpital Civil - BP 246 - 67091 STRASBOURG CEDEX - FRANCE

In medical professions, identification and evaluation of health: disorders are more difficult than in the general population. Making available self-tests on mental health, behavioral disruptions, abusive consumptions and sleep disorders could help doctors and nurses to be aware of their own health. The French college of anesthetists (CFAR) provides free access to these tests on its website since October 2013 (www.cfar.org). Analysis of connections and completions are presented.

METHOD AND RESULTS

Nine internationally validated self-assessment tests (8) internationally validated, and an additional test on harassment [validation in progress]) were made available on line for anesthetist nurses and physicians (7 since October 2013 and 2, harassment and workaholism, in summer 2014). Connections and full completion were analysed. Anonymity was fully preserved and no information on result details was possible.

Available self-assessments on www.cfar.org website	Connected	Completed	%Completed /Connected	Ranking by completing frequency
Burn-out (Maslach Burnout Inventory)	868	811	93%	26%
Anxiety-depression (Hospital Anxiety Depression sc.)	618	476	77%	16%
Drowsiness (EPWORTH)	535	470	88%	15%
Fatigue (PICHOT)	595	441	74%	14%
Alcohol (FACE)	452	370	82%	12%
Cyber addiction	354	207	58%	7%
Tobacco (FAGERSTRÖM)	261	93	36%	3%
Harassment	250	103	41%	3%
Workalholism (Bergen Work Addiction Scale	146	98	67%	3%
TOTAL	4079	3069	75%	100%
CFAR website www.cfar.or	g / SMART ta	b. (APRIL-06-2	2015)	

Downloading anonymous report form is possible for personal data.

- 1. This also allows individual, or better, global analysis in the aim to obtain a team evaluation.
- 2. Eventually, when abnormal results occur, the team can clearly make organizational changes.

	In case of collective evaluation, don't forg by the Project Manager in your team :;		· · · · · · · · · · · · · · · · · · ·		
CFAR	« Health – Work – Perso	onal Li	ife » Assessment		
SMART VIE PR	confidentially HAD tests, Burnout, Patigue and ATIQUE / AUTOTESTS) or (http://www.cfar.org				
Position:	☐ Intern ☐ Resident ☐ Assistant ☐ Cons	altant 🗆 .	Acad. 🗆 Private 🗆 prefer to stay di		
□ (précisez :	Murse Anesth OR Nurse Nurse 	Health M	anager 🔲 Other HC Provider		
Seniority i	this HealthCare Institution : \square < 5 ye	ers	☐ 5 à 10 years ☐ > 10 years		
Test HAD (Hospital Anxiety – Depression Scale)				
	icore Anxiety :/21 icore Depression:/21	810	Interpretation 8 to 10 : anxiety or depression are unlikely More than 10 : anxiety or depression proved		
Burnout (N	ABI — Maslach Burnout Inventory)				
-	motional Exhaustion(EE) score/54 epersonalization (DP) score/30	1	CORRECT ATTENTION DANGEROUS 0 18 30 Otyperson nativation CONFECT ATTENTION DANGEROUS 0 6 12		
• My P	ersonal Accomplishment(AP) / 48		0 34 40		
Fatigue (Pi	chot's Test)				
 My Score 	: Pichot:/ 32				
Drowsines	s (Epworth)	45	No sleep debt		
My score Epworth's Scale :/24		9 à 14	The state of the s		
_	•	h 15	You experience excessive daytime alsopiness sig		
-	eekly hours devoted to work				
	of my presence in the Institution can be e at working at Home for Institution (Record to		-		
•		AND DESCRIPTIONS	ancerome, at		
	lumber:Children				
Optional: A	ge rating: 🗆 20-25 🗎 26-35 🗀 36-45 🗆 4	16-55 S	6-65 □ >65		
Addiction to v identified that specialist - tol	ther tests are available on the CFAR website (To vork). They do not base a diagnosis but allow av t could justify making contact with a profession racco specialist, psychologist,	vareness to al to take s	o whom permits. So, conditions could l tock: medical workers, GP, addiction		
organization r success in this	n, an anonymous global assessment with all the ecognized as unsatisfactory or pathogens. The approach. In this case, anonymously complete	participatio	on of all team members is a key factor		
(leader, CDS).	To know more:www.cfar.org		s www.cfar.org Thanks for particip		

> DISCUSSION

- > Up to 6th April 2015, there have been 4079 connections on self-tests, 3069 of which: were fully completed, corresponding to 1/3 of the target population of French: Anesthetists.
- > MBI test, Anxiety-Depression, Fatigue, Epworth and Alcohol, were the tests the most frequently selected (21%, 15%, 15%, 13%, 11% respectively) and fully completed (93%, 77%, : 74%, 88%, 82% respectively).
- > Taking into account the time needed to complete on line the harassment and workaholism tests, results suggest that the tests are consulted but less fully completed: (harassment test), due to the length of time required to complete them for exhaustive and informative data.
- > Results suggest that information on their own health state can be self-assessed by anaesthesiologists and intensivists in France. Particularly concerning mental or exhaustion disorders and addiction however less for tobacco dependency.

> CONCLUSIONS

Confronted with denial, the use of such tests could aid doctors to initiate a therapeutic action with a variety of specialists: own addiction specialist, doctor, chosen psychiatrist. However psychologist or systemic approach is possible when team affected members organizational dysfunction exhaustion, over-workload, lack of resources, inequity, isolation, harassment...

www.cfar.org









